Available online at www.sciencedirect.com
SCIENCE DIRECT®
ELSEVIER
Aquaculture 221 (2003) 667–670
Aquaculture
www.elsevier.com/locate/aqua-online

Contents of *Aquaculture*, Volume 221

VOL. 221 NOS. 1–4 **1 May 2003**

Diseases, Parasites and Predators

Biochemical, physiological and hematological changes in white spot syndrome virus-infected shrimp, *Penaeus indicus*
K. Yoganandhan (Chennai, India), S. Thirupathi and A.S. Sahul Hameed (Vellore, India) 1

Effects of copper-based antifouling treatment on the presence of *Neoparamoeba pemaquidensis* Page, 1987 on nets and gills of reared Atlantic salmon (*Salmo salar*)
G.M. Douglas-Helders, C. Tan, J. Carson and B.F. Nowak (Launceston, Tas., Australia) 13

Expression of immune genes in turbot (*Scophthalmus maximus*) fed a nucleotide-supplemented diet
C. Low (Aberdeen, UK), S. Wadsworth (Argyll, UK), C. Burrells (Dirdal, Norway) and C.J. Secombes (Aberdeen, UK) 23

Hematological and biochemical parameters in common carp, *Cyprinus carpio*, following herbal treatment for *Aeromonas hydrophila* infection
R. Harikrishnan, M. Nisha Rani and C. Balasundaram (Tiruchirapalli, India) 41

Effects of oral treatment with TNP-470 on rainbow trout (*Oncorhynchus mykiss*) infected with *Tetracapsuloides bryosalmonae* (Malacosporea), the causative agent of proliferative kidney disease
D.J. Morris, A. Adams (Stirling, UK), P. Smith (Saffron Walden, UK) and R.H. Richards (Stirling, UK) 51

Isolation of *Vibrio tapetis* from cultivated Atlantic halibut (*Hippoglossus hippoglossus* L.)
H.I. Reid, H.L. Duncan (Glasgow, UK), L.A. Laidler, D. Hunter (Inverness-shire, UK) and T.H. Birkbeck (Glasgow, UK) 65

Pharmacokinetics and bioavailability of oxytetracycline in gilthead sea bream (*Sparus aurata*) after a single dose
G. Rigos (Attiki, Greece; Kingston upon Thames, UK), I. Nengas (Attiki, Greece), A.E. Tyrpenou (Athens, Greece), M. Alexis (Attiki, Greece) and G.M. Troisi (Middlesex, UK) 75

Spionid mudworm infestation of farmed abalone (*Haliotis* spp.)
M. Lleonart (King Meadows, Launceston, Tas., Australia), J. Handlinger (King Meadows, Tas., Australia) and M. Powell (Launceston, Tas., Australia) 85

Histopathology of Y-organ in Indian white shrimp *Fenneropenaeus indicus*, experimentally infected with white spot syndrome virus
K.K. Vijayan, C.P. Balasubramanian, K.P. Jithendran, S.V. Alavandi and T.C. Santiago (Chennai, India) 97

Differences between short- and long-term cultures of *Uronema marinum* (Ciliophora: Scuticociliatida) in chemiluminescence inhibitory activity, antioxidative enzyme and protease activity
S.R. Kwon, C.S. Kim and K.H. Kim (Pusan, South Korea) 107

Genetics

The competitiveness of the eel aquaculture in Taiwan, Japan, and China
W.-C. Lee (Taipei, Keelung, Taiwan), Y.-H. Chen, Y.-C. Lee (Taipei, Taiwan) and I.C. Liao (Keelung, Taiwan) 115

An investigation of sex determination in the Mozambique tilapia, *Oreochromis mossambicus*, using synaptonemal complex analysis, FISH, sex reversal and gynogenesis
R. Campos-Ramos, S.C. Harvey, B.J. McAndrew and D.J. Penman (Stirling, UK) 125

Husbandry and Management

Effects of temperature on initial swim bladder inflation and related development in cultured striped trumpeter (*Latris lineata*) larvae
A.J. Trotter (Launceston, Hobart, Tas., Australia), P.M. Pankhurst (Launceston, Tas., Australia), D.T. Morehead and S.C. Battaglene (Hobart, Tas., Australia) 141

The use of ozone in trout hatchery to reduce saprolegniasis incidence
G. Forneris, S. Bellardi, G.B. Palmegiano (Grugliasco, Italy), M. Saroglia (Varese, Italy), B. Sicuro, L. Gasco and I. Zoccarato (Grugliasco, Italy) 157

A semi-recirculating, integrated system for the culture of fish and seaweed
A. Schuenhoff (Faro, Portugal), M. Shpigel, I. Lupatsch, A. Ashkenazi, F.E. Msuya and A. Neori (Eilat, Israel) 167

Effects of rearing density, size grading and parental factors on sex ratios of the sea bass (*Dicentrarchus labrax* L.) in intensive aquaculture
E. Saillant (Palavas-les-Flots, Gravelines, France), A. Fostier, P. Haffray (Rennes, France), B. Menu (Palavas-les-Flots, France), S. Laureau (Gravelines, France), J. Thimonier (Montpellier, France) and B. Chatain (Palavas-les-Flots, France) 183

Evaluation of biological and economical aspects of the culture of the scallop *Lyropecten (Nodipecten) nodosus* in suspended and bottom culture
Y. Mendoza, L. Freites, C.J. Lodeiros, J.A. López (Cumaná, Venezuela) and J.H. Himmelman (Quebec, Canada) 207

Culture-based fisheries: an underutilised opportunity in aquaculture development
S.S. De Silva (Warrnambool, Australia) 221

Hatchability of eggs from Atlantic cod, turbot and Atlantic halibut after disinfection with ozonated seawater
S. Grotmol, E. Dahl-Paulsen and G.K. Totland (Bergen, Norway) 245

Microsatellite-based pedigree tracing in a Japanese flounder *Paralichthys olivaceus* hatchery strain: implications for hatchery management related to stock enhancement program
M. Sekino, K. Saitoh (Shiogama, Japan), T. Yamada, A. Kumagai (Miyako, Japan), M. Hara (Watarai, Japan) and Y. Yamashita (Maizuru, Japan) 255

Advanced nursing of mixed-sex and mono-sex tilapia (*Oreochromis niloticus*) fry, and its impact on subsequent growth in fertilized ponds
D.C. Little (Pathumthani, Thailand; Scotland, UK), R.C. Bhujel (Pathumthani, Thailand) and T.A. Pham (Bac Ninh, Viet Nam) 265

Growth and survival of pacu *Piaractus mesopotamicus* (Holmberg, 1887) juveniles reared in ponds or at different initial larviculture periods indoors
R.K. Jomori, D.J. Carneiro, E.B. Malheiros (São Paulo, Brazil) and M.C. Portella (Brazil) 277

The toxicity of emamectin benzoate, an aquaculture pesticide, to planktonic marine copepods
K.J. Willis (Oban, UK) and N. Ling (Hamilton, New Zealand) 289

Interpreting evidence of dispersal by *Haliotis midae* juveniles seeded in the wild
S.W.P. de Waal, G.M. Branch and R. Navarro (Cape Town, South Africa) 299

Sponge aquaculture for the production of biologically active metabolites: the influence of farming protocols and environment
A. Duckworth and C. Battershill (Wellington, New Zealand) 311

Relationship between physicochemical variables and productivity in open ponds for the production of *Spirulina*: a predictive model of algal yield
C. Jiménez, B.R. Cossío and F.X. Niell (Málaga, Spain) 331

The effect of co-cultivation on the pearl yield of *Pinctada martensi* (Dumker)
M. Wu (Hong Kong, China), S.K.K. Mak (Hong Kong, China), X. Zhang (China) and P.-Y. Qian (Hong Kong, China) 347

Evaluation of mucus, *Navicula*, and mixed diatoms as larval settlement inducers for the tropical abalone *Haliotis asinina*
W.G. Gallardo and S.M.A. Buen (Iloilo, Philippines) 357

Nutrition

Efficiency of feed utilization in Atlantic salmon (*Salmo salar* L.) fed diets with increasing substitution of fish meal with vegetable proteins
J. Opstvedt, A. Aksnes, B. Hope (Bergen, Norway) and I.H. Pike (St. Albans, UK) 365
A retinoic acid receptor-selective agonist causes jaw deformity in the Japanese flounder, *Paralichthys olivaceus*
Y. Haga (Tokyo, Japan), T. Suzuki (Mie, Japan), H. Kagechika and T. Takeuchi (Tokyo, Japan) . . . 381
Variation of lipid class and fatty acid composition of *Chaetoceros muelleri* and *Isochrysis* sp. grown in a semicontinuous system
F. Pernet, R. Tremblay (Quebec, Canada), E. Demers (Gaspe, Quebec, Canada) and M. Roussy (Quebec, Canada) 393
Anthropogenic effects on seston quality and quantity and the growth and survival of Sydney rock oyster (*Saccostrea glomerata*) in two estuaries in NSW, Australia
K.J. Paterson, M.J. Schreider and K.D. Zimmerman (Ourimbah, Australia) 407
Utilization of glucose, maltose, dextrin and cellulose by juvenile flounder (*Paralichthys olivaceus*)
S.-M. Lee, K.-D. Kim (Kangnung, South Korea) and S.P. Lall (Halifax, NS, Canada) 427
Growth and feed conversion of juvenile winter flounder (*Pleuronectes americanus*) in relation to different protein-to-lipid levels in isocaloric diets
C.D. Hebb (Truro, NS, Canada), J.D. Castell (St. Andrew's, NB, Canada), D.M. Anderson (Truro, NS, Canada) and J. Batt (Halifax, NS, Canada) 439
Dietary phosphorus requirement of juvenile haddock (*Melanogrammus aeglefinus* L.)
P.K. Roy and S.P. Lall (Halifax, NS, Canada) 451
Modulation of glucose-6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase and malic enzyme activity by glucose and alanine in Atlantic salmon, *Salmo salar* L. hepatocytes
M. Sanden, L. Frøyland and G.-I. Hemre (Bergen, Norway) 469
An acoustic detector of turbot feeding activity
R. Mallekh, J.P. Lagardère (L'Houmeau, France), J.P. Eneau (Nantes, France) and C. Cloutour (Noirmoutier, France) 481
Nutritional value of fresh and concentrated algal diets for larval and juvenile Pacific oysters (*Crassostrea gigas*)
E. Ponis (Florence, Italy), R. Robert (Landunvez, France) and G. Parisi (Florence, Italy) 491
Biochemical changes during the embryonic development of Norway lobster, *Nephrops norvegicus*
R. Rosa (Lisbon, Portugal), S. Morais, R. Calado, L. Narciso (Cascais, Portugal) and M.L. Nunes Lisbon, Portugal) 507

Physiology and Endocrinology

Improved reproductive readiness of pond-reared broodstock *Penaeus merguiensis* by environmental manipulation
T. Hoang, S.Y. Lee, C.P. Keenan (Queensland, Australia) and G.E. Marsden (Grafton, NSW, Australia) 523
Impairment of thymus development in cultured osmerid fish, the ayu, *Plecoglossus altivelis*
S. Miwa (Yokohama, Japan), A. Sakai (Hikone, Japan) and M. Nakane (Yokohama, Japan) 535
The efficacy of metomidate, clove oil, Aqui-S™ and Benzoak® as anaesthetics in Atlantic salmon (*Salmo salar* L.) smolts, and their potential stress-reducing capacity
M. Iversen (Trondheim, Bodö, Norway), B. Finstad (Trondheim, Norway), R.S. McKinley (West Vancouver, Canada) and R.A. Eliassen (Bodö, Norway) 549
Long-to-short photoperiod in winter halves the incidence of sexual maturity among Arctic charr
J. Duston, T. Astatkie and P.F. MacIsaac (Truro, Nova Scotia, Canada) 567

Artificial photoperiod influence on the immune system of juvenile rainbow trout (*Oncorhynchus mykiss*) in the Southern Hemisphere
M.O. Leonardi and A.E. Klempau (Concepción, Chile) 581
Effect of different treatments on the chorion permeability to DMSO of turbot embryos (*Scophthalmus maximus*)
E. Cabrita, O. Chereguini, M. Luna, P. de Paz and M.P. Herráez (Leon, Spain). 593
Multiple spawning and egg quality of individual European sea bass (*Dicentrarchus labrax*) females after repeated injections of GnRHa
C.C. Mylonas, I. Sigelaki, P. Divanach (Crete, Greece), E. Mananõs, M. Carrillo (Castellón, Spain) and A. Afonso-Polyviou (Nicosia, Cyprus). 605
Effects of dietary conjugated linoleic acids and total dietary lipid concentrations on growth responses of juvenile channel catfish, *Ictalurus punctatus*
R.G. Twibell and R.P. Wilson (Mississippi State, MS, USA). 621
Physiology and survival of triploid brook trout following exhaustive exercise in warm water
C.A. Hyndman, J.D. Kieffer (Saint John, NB, Canada) and T.J. Benfey (Fredericton, NB, Canada) . 629
Effect of ovarian stage on plasma vitellogenin and calcium in cultured white sturgeon
J. Linares-Casenave, K.J. Kroll, J.P. Van Eenennaam and S.I. Doroshov (Davis, CA, USA) 645
Residue depletion of oxytetracycline from fillet tissues of northern pike and walleye
J.A. Bernardy, C. Vue, M.P. Gaikowski, G.R. Stehly, W.H. Gingerich (Crosse, WI, USA) and A. Moore (Moravia, IA, USA) 657

Contents of *Aquaculture*, Volume 221 667

Contents (continued)

Effects of oral treatment with TNP-470 on rainbow trout (*Oncorhynchus mykiss*) infected with *Tetracapsuloides bryosalmonae* (Malacosporea), the causative agent of proliferative kidney disease
D.J. Morris, A. Adams (Stirling, UK), P. Smith (Saffron Walden, UK) and R.H. Richards (Stirling, UK) 51
Isolation of *Vibrio tapetis* from cultivated Atlantic halibut (*Hippoglossus hippoglossus* L.)
H.I. Reid, H.L. Duncan (Glasgow, UK), L.A. Laidler, D. Hunter (Inverness-shire, UK) and T.H. Birkbeck (Glasgow, UK) 65
Pharmacokinetics and bioavailability of oxytetracycline in gilthead sea bream (*Sparus aurata*) after a single dose
G. Rigos (Attiki, Greece; Kingston upon Thames, UK), I. Nengas (Attiki, Greece), A.E. Tyrpenou (Athens, Greece), M. Alexis (Attiki, Greece) and G.M. Troisi (Middlesex, UK) 75
Spionid mudworm infestation of farmed abalone (*Haliotis* spp.)
M. Lleonart (King Meadows, Launceston, Tas., Australia), J. Handlinger (King Meadows, Tas., Australia) and M. Powell (Launceston, Tas., Australia) 85
Histopathology of Y-organ in Indian white shrimp *Fenneropenaeus indicus*, experimentally infected with white spot syndrome virus
K.K. Vijayan, C.P. Balasubramanian, K.P. Jithendran, S.V. Alavandi and T.C. Santiago (Chennai, India) 97
Differences between short- and long-term cultures of *Uronema marinum* (Ciliophora: Scuticociliatida) in chemiluminescence inhibitory activity, antioxidative enzyme and protease activity
S.R. Kwon, C.S. Kim and K.H. Kim (Pusan, South Korea) 107

Genetics

The competitiveness of the eel aquaculture in Taiwan, Japan, and China
W.-C. Lee (Taipei, Keelung, Taiwan), Y.-H. Chen, Y.-C. Lee (Taipei, Taiwan) and I.C. Liao (Keelung, Taiwan) 115
An investigation of sex determination in the Mozambique tilapia, *Oreochromis mossambicus*, using synaptonemal complex analysis, FISH, sex reversal and gynogenesis
R. Campos-Ramos, S.C. Harvey, B.J. McAndrew and D.J. Penman (Stirling, UK) 125

Husbandry and Management

Effects of temperature on initial swim bladder inflation and related development in cultured striped trumpeter (*Latris lineata*) larvae
A.J. Trotter (Launceston, Hobart, Tas., Australia), P.M. Pankhurst (Launceston, Tas., Australia), D.T. Morehead and S.C. Battaglene (Hobart, Tas., Australia) 141
The use of ozone in trout hatchery to reduce saprolegniasis incidence
G. Forneris, S. Bellardi, G.B. Palmegiano (Grugliasco, Italy), M. Saroglia (Varese, Italy), B. Sicuro, L. Gasco and I. Zoccarato (Grugliasco, Italy) 157
A semi-recirculating, integrated system for the culture of fish and seaweed
A. Schuenhoff (Faro, Portugal), M. Shpigel, I. Lupatsch, A. Ashkenazi, F.E. Msuya and A. Neori (Eilat, Israel) 167
Effects of rearing density, size grading and parental factors on sex ratios of the sea bass (*Dicentrarchus labrax* L.) in intensive aquaculture
E. Saillant (Palavas-les-Flots, Gravelines, France), A. Fostier, P. Haffray (Rennes, France), B. Menu (Palavas-les-Flots, France), S. Laureau (Gravelines, France), J. Thimonier (Montpellier, France) and B. Chatain (Palavas-les-Flots, France) 183

Contents (continued)

Evaluation of biological and economical aspects of the culture of the scallop *Lyropecten (Nodipecten) nodosus* in suspended and bottom culture
Y. Mendoza, L. Freites, C.J. Lodeiros, J.A. López (Cumaná, Venezuela) and J.H. Himmelman (Quebec, Canada) 207
Culture-based fisheries: an underutilised opportunity in aquaculture development
S.S. De Silva (Warrnambool, Australia) 221
Hatchability of eggs from Atlantic cod, turbot and Atlantic halibut after disinfection with ozonated seawater
S. Grotmol, E. Dahl-Paulsen and G.K. Totland (Bergen, Norway). 245
Microsatellite-based pedigree tracing in a Japanese flounder *Paralichthys olivaceus* hatchery strain: implications for hatchery management related to stock enhancement program
M. Sekino, K. Saitoh (Shiogama, Japan), T. Yamada, A. Kumagai (Miyako, Japan), M. Hara (Watarai, Japan) and Y. Yamashita (Maizuru, Japan). 255
Advanced nursing of mixed-sex and mọno-sex tilapia (*Oreochromis niloticus*) fry, and its impact on subsequent growth in fertilized ponds
D.C. Little (Pathumthani, Thailand; Scotland, UK), R.C. Bhujel (Pathumthani, Thailand) and T.A. Pham (Bac Ninh, Viet Nam). 265
Growth and survival of pacu *Piaractus mesopotamicus* (Holmberg, 1887) juveniles reared in ponds or at different initial larviculture periods indoors
R.K. Jomori, D.J. Carneiro, E.B. Malheiros (São Paulo, Brazil) and M.C. Portella (Brazil). . 277
The toxicity of emamectin benzoate, an aquaculture pesticide, to planktonic marine copepods
K.J. Willis (Oban, UK) and N. Ling (Hamilton, New Zealand) 289
Interpreting evidence of dispersal by *Haliotis midae* juveniles seeded in the wild
S.W.P. de Waal, G.M. Branch and R. Navarro (Cape Town, South Africa) 299
Sponge aquaculture for the production of biologically active metabolites: the influence of farming protocols and environment
A. Duckworth and C. Battershill (Wellington, New Zealand) 311
Relationship between physicochemical variables and productivity in open ponds for the production of *Spirulina*: a predictive model of algal yield
C. Jiménez, B.R. Cossío and F.X. Niell (Málaga, Spain) 331
The effect of co-cultivation on the pearl yield of *Pinctada martensi* (Dumker)
M. Wu (Hong Kong, China), S.K.K. Mak (Hong Kong, China), X. Zhang (China) and P.-Y. Qian (Hong Kong, China) 347
Evaluation of mucus, *Navicula*, and mixed diatoms as larval settlement inducers for the tropical abalone *Haliotis asinina*
W.G. Gallardo and S.M.A. Buen (Iloilo, Philippines) 357

Nutrition

Efficiency of feed utilization in Atlantic salmon (*Salmo salar* L.) fed diets with increasing substitution of fish meal with vegetable proteins
J. Opstvedt, A. Aksnes, B. Hope (Bergen, Norway) and I.H. Pike (St. Albans, UK) 365
A retinoic acid receptor-selective agonist causes jaw deformity in the Japanese flounder, *Paralichthys olivaceus*
Y. Haga (Tokyo, Japan), T. Suzuki (Mie, Japan), H. Kagechika and T. Takeuchi (Tokyo, Japan). 381
Variation of lipid class and fatty acid composition of *Chaetoceros muelleri* and *Isochrysis* sp. grown in a semicontinuous system
F. Pernet, R. Tremblay (Quebec, Canada), E. Demers (Gaspe, Quebec, Canada) and M. Roussy (Quebec, Canada). 393

Contents (continued)

Anthropogenic effects on seston quality and quantity and the growth and survival of Sydney rock oyster (*Saccostrea glomerata*) in two estuaries in NSW, Australia
K.J. Paterson, M.J. Schreider and K.D. Zimmerman (Ourimbah, Australia) 407
Utilization of glucose, maltose, dextrin and cellulose by juvenile flounder (*Paralichthys olivaceus*)
S.-M. Lee, K.-D. Kim (Kangnung, South Korea) and S.P. Lall (Halifax, NS, Canada) 427
Growth and feed conversion of juvenile winter flounder (*Pleuronectes americanus*) in relation to different protein-to-lipid levels in isocaloric diets
C.D. Hebb (Truro, NS, Canada), J.D. Castell (St. Andrew's, NB, Canada), D.M. Anderson (Truro, NS, Canada) and J. Batt (Halifax, NS, Canada) 439
Dietary phosphorus requirement of juvenile haddock (*Melanogrammus aeglefinus* L.)
P.K. Roy and S.P. Lall (Halifax, NS, Canada) 451
Modulation of glucose-6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase and malic enzyme activity by glucose and alanine in Atlantic salmon, *Salmo salar* L. hepatocytes
M. Sanden, L. Frøyland and G.-I. Hemre (Bergen, Norway) 469
An acoustic detector of turbot feeding activity
R. Mallekh, J.P. Lagardère (L'Houmeau, France), J.P. Eneau (Nantes, France) and C. Cloutour (Noirmoutier, France) 481
Nutritional value of fresh and concentrated algal diets for larval and juvenile Pacific oysters (*Crassostrea gigas*)
E. Ponis (Florence, Italy), R. Robert (Landunvez, France) and G. Parisi (Florence, Italy) . . 491
Biochemical changes during the embryonic development of Norway lobster, *Nephrops norvegicus*
R. Rosa (Lisbon, Portugal), S. Morais, R. Calado, L. Narciso (Cascais, Portugal) and M.L. Nunes (Lisbon, Portugal) 507

Physiology and Endocrinology

Improved reproductive readiness of pond-reared broodstock *Penaeus merguiensis* by environmental manipulation
T. Hoang, S.Y. Lee, C.P. Keenan (Queensland, Australia) and G.E. Marsden (Grafton, NSW, Australia) 523
Impairment of thymus development in cultured osmerid fish, the ayu, *Plecoglossus altivelis*
S. Miwa (Yokohama, Japan), A. Sakai (Hikone, Japan) and M. Nakane (Yokohama, Japan) 535
The efficacy of metomidate, clove oil, Aqui-S™ and Benzoak® as anaesthetics in Atlantic salmon (*Salmo salar* L.) smolts, and their potential stress-reducing capacity
M. Iversen (Trondheim, Bodö, Norway), B. Finstad (Trondheim, Norway), R.S. McKinley (West Vancouver, Canada) and R.A. Eliassen (Bodö, Norway) 549
Long-to-short photoperiod in winter halves the incidence of sexual maturity among Arctic charr
J. Duston, T. Astatkie and P.F. MacIsaac (Truro, Nova Scotia, Canada) 567
Artificial photoperiod influence on the immune system of juvenile rainbow trout (*Oncorhynchus mykiss*) in the Southern Hemisphere
M.O. Leonardi and A.E. Klempau (Concepción, Chile) 581
Effect of different treatments on the chorion permeability to DMSO of turbot embryos (*Scophthalmus maximus*)
E. Cabrita, O. Chereguini, M. Luna, P. de Paz and M.P. Herráez (Leon, Spain) 593
Multiple spawning and egg quality of individual European sea bass (*Dicentrarchus labrax*) females after repeated injections of GnRHa
C.C. Mylonas, I. Sigelaki, P. Divanach (Crete, Greece), E. Mananõs, M. Carrillo (Castellón, Spain) and A. Afonso-Polyviou (Nicosia, Cyprus) 605

Contents (continued)

Effects of dietary conjugated linoleic acids and total dietary lipid concentrations on growth responses of juvenile channel catfish, *Ictalurus punctatus*
R.G. Twibell and R.P. Wilson (Mississippi State, MS, USA) 621
Physiology and survival of triploid brook trout following exhaustive exercise in warm water
C.A. Hyndman, J.D. Kieffer (Saint John, NB, Canada) and T.J. Benfey (Fredericton, NB, Canada) 629
Effect of ovarian stage on plasma vitellogenin and calcium in cultured white sturgeon
J. Linares-Casenave, K.J. Kroll, J.P. Van Eenennaam and S.I. Doroshov (Davis, CA, USA) 645
Residue depletion of oxytetracycline from fillet tissues of northern pike and walleye
J.A. Bernardy, C. Vue, M.P. Gaikowski, G.R. Stehly, W.H. Gingerich (Crosse, WI, USA) and A. Moore (Moravia, IA, USA) 657

Contents of *Aquaculture*, Volume 221 667